



MARSH MODEL RETROSPECTIVE WORKSHOP

APRIL 11 -12, 2022



Authors: Sara Martin, Charley Cameron, Renee Collini, Nina Woodard, Christine Buckel, Trevor Meckley, Sarah Spiegler, Karim Alizad, Harris Bienn, Jonathan Clough, Christopher Esposito, Stephen Medeiros, James Morris, Mary Schoell

MASGP-22-055

The project described in this publication was supported by the U.S. Department of Commerce's National Oceanic and Atmospheric Administration under NOAA Award NA18OAR4170080 and the Mississippi-Alabama Sea Grant Consortium. Its contents are solely the responsibility of the authors and do not necessarily represent the views of the USGS Southeast Climate Adaptation Science Center or the USGS. This manuscript is submitted for publication with the understanding that the United States Government is authorized to reproduce and distribute reprints for Governmental purposes.

Table of Contents

List of Figures and Tables.....	5
List of Acronyms.....	6
1.0 Introduction	6
1.1 Objectives	8
2.0 Attendee List	9
3.0 Discussion Summaries	10
3.1 Stakeholder Perceptions on Marsh Modeling	10
3.1.1 Decision Points	11
3.1.2 Next Steps	11
3.2 Big Picture Questions for the Marsh Model Retrospective	11
3.3 Available Data for Performing a Marsh Model Retrospective	12
3.3.1 Decision Points	14
3.3.2 Next Steps	14
3.4 Day One Wrap-Up	14
3.5 Details for Performing a Marsh Model Retrospective	14
3.5.1 Vegetation Input Layers	14
3.5.1.1 Next Steps	15
3.5.2 Digital Elevation Models.....	16
3.5.2.1 Next Steps	16
3.5.3 Time Steps.....	16
3.5.3.1 Next Steps	17
3.5.4 Sea-Level Rise.....	17
3.5.4.1 Next Steps	17
3.5.5 Locations.....	17
3.5.5.1 Grand Bay NERR.....	18
3.5.5.2 Apalachicola NERR	18
3.5.5.3 North Carolina Reserve System	18
3.5.5.4 Louisiana.....	18
3.5.5.5 Plum Island, MA.....	18
3.5.5.6 Sacramento Delta	18
3.5.5.7 Next Steps	19
3.5.6 Model Uncertainty	19
3.5.6.1 Next Steps	20
3.6 Discussion Summary: Revisiting the Marsh Model Retrospective Framework	20
3.7 An Approach and Timeline for the Marsh Model Retrospective	21
3.7.1 Next Steps	23
4.0 Appendices	24
Appendix A: Workshop Presentations	25
A1: Introduction to the Workshop	25
A2: Stakeholder Perceptions of Marsh Models	27
A3: Background on the Marsh Model Retrospective.....	40
A4: Introduction to Data Exploration	43
A5: Marsh Model Retrospective Framework	61
Appendix B: Handouts	64
B1: Participant Agenda	64
B2: Questions for Marsh Model Retrospective	66

B3: Data Inventory.....	67
B4: Map Data Exploration Activity Sheet	75
Appendix C: Data Exploration Web Application Screenshots	76
Appendix D: Workshop Flip Charts.....	78
D1: Workshop Marina & Ground Rules.....	78
D2: Managers Want Modelers to Know	79
D3: Things for Managers to Know.....	80
D4: Data Discussion.....	81
D5: Details of the Retrospective.....	82
D6: Scoping Question Discussion.....	83
D7: Next Steps	83

List of Figures and Tables

Figure 1: Screenshot of the web mapping platform used for the data exploration by the marsh shoreline

12

Table 1. List of models represented in the workshop. 6

Table 2. Managers Want Models That... 9

Table 3. Things for Managers to Know 9

Table 4. Considerations for Scoping the Marsh Model Retrospective 11

Table 5. Data Concerns 12

Table 6. Data Needs 12

Table 7. Data Want 13

Table 8. Marsh Model Retrospective Outputs to Compare 13

Table 9. Model Vegetation Data Definitions and Other Important Data 14

Table 10. Votes for Marsh Model Retrospective Locations 18

Table 11. Sources of Uncertainty in Models 19

List of Acronyms

CCAP	Coastal Change Analysis Program
DEM	Digital Elevation Model
Hydro-MEM	Hydrodynamic-Marsh Equilibrium Model
ICM	Integrated Compartment Model
MEM	Marsh Equilibrium Model
NERR	National Estuarine Research Reserve
NOAA	National Oceanic and Atmospheric Administration
NWI	National Wetland Inventory
PLACE:SLR	Program For Local Adaptation to Climate Effects: Sea Level Rise
RTK	Real Time Kinematic
SC	South Carolina
SET	Surface Elevation Table
SLAMM	Sea Level Affecting Marshes Model
TSS	Total Suspended Sediments
USGS	United States Geological Survey
WARMER	Wetland Accretion Rate Model of Ecosystem Resilience

1.0 Introduction

On August 7-9, 2018, natural resource managers and decision makers from across the Gulf of Mexico were convened to explore comparisons of already existing marsh model outputs and discuss potential drivers of the differences and how this may impact choices when selecting models to support natural resource decision making. The purpose of this comparison was not to identify the “best” model, but to instead work on understanding the different outputs of each model and how managers might utilize the different marsh models for different purposes. At this workshop, there was agreement amongst the participants that a retrospective analysis needed to be performed with all the models utilizing the same data inputs across multiple geographies. The results of the retrospective analysis would enhance guidelines on the model application and identify potential areas of research to enhance the existing models’ predictive capabilities.

A Project Team comprised of marsh model funders, coastal resilience specialists who support coastal managers, data analysis and visualization specialists, and modeling experts secured funding to convene marsh modelers from a range of marsh models in order to scope a retrospective analysis. On April 11–12, 2022, marsh modelers representing six marsh models (Table 1) gathered with the Project Team to discuss an approach and timeline for a marsh model retrospective at the Beaufort Hotel in Beaufort, North Carolina. This workshop had six objectives that were achieved throughout the course of the two days.

Table 1. List of marsh models and the representative in attendance at the April 11-12, 2022 workshop.

Marsh Model Name	Model Representative
Hydro-MEM	Karim Alizad
Integrated Compartment Model (ICM)	Eric White
Marsh Equilibrium Model (MEM)	James Morris
Sea Level Affecting Marshes Model (SLAMM)	Jonathan Clough
Sea Level Rise Viewer (SLR Viewer)	Connor Levy
Wetland Accretion Rate Model of Ecosystem Resilience (WARMER)	Kevin Buffington

The first day of the workshop (April 11) consisted of presentations, discussions, and activities developed and led by the Project Team to prepare the marsh modelers for in depth discussion of the technical details required to perform a marsh model retrospective. These introductory sessions were conducted to accomplish the following objectives. To:

1. learn about stakeholder perceptions of marsh models to ensure that the work resulting from the workshop is informed by the needs expressed by stakeholders;
2. reach a consensus about what a marsh model retrospective would look like and be able to accomplish; and
3. explore the historical data that are available for performing a marsh model retrospective analysis.

Through activities and robust discussions, each of these objectives were accomplished during day one of the workshop. See sections 3.1-3.4 for day one discussion summaries.

The overarching goal of day two (April 12) of the workshop was to determine sufficient technical details to develop a proposal to advance the marsh model retrospective project. This was accomplished through detailed discussions that shared both the knowledge that the Project Team had accumulated in preparation for the workshop and the professional expertise of the marsh modelers, in addition to collaboration between both groups for future planning. During day two, workshop participants (the Project Team and the modelers in collaboration) accomplished the following objectives:

1. selected locations and other technical details such as how to develop historic DEMs, estimate sea-level rise over the period of record of the retrospective, etc.;
2. determined the next steps to assess and prepare the model input data; and
3. agreed upon a timeline for completing the next steps for performing the marsh model retrospective analysis.

See sections 3.5-3.7 for day two discussion summaries.

By the end of the workshop, the marsh modelers understood how stakeholders perceive marsh models, the need for and goals of a marsh model retrospective, what data are available for performing a marsh model retrospective, and where additional data gaps may lie. This laid the foundation for the marsh modelers to collaborate with the Project Team to complete a framework for performing the marsh model retrospective, including what parameters to include, next steps, and a timeline for completing the prerequisite work for the marsh model retrospective. Overall, this two-day workshop successfully prepared the Project Team and marsh modelers to gather necessary data and move the marsh model retrospective forward.

1.1 Objectives

Day One (April 11, 2022)

- The marsh modelers will understand how stakeholders use and perceive marsh model outputs in their decision-making.
- The project team and marsh modelers will agree upon the scope of the marsh model retrospective and understand what it will accomplish.
- The marsh modelers will have a better understanding of what historical data is available for performing a marsh model retrospective.

Day Two (April 12, 2022)

- The marsh modelers and project team will determine the priority locations and select other technical parameters for performing the marsh model retrospective analysis.
- The marsh modelers and project team will collaborate to develop next steps and assign responsibilities for the next steps to move the marsh model retrospective analysis forward.
- The marsh modelers and project team will develop and agree upon a timeline for all next steps for the marsh model retrospective analysis.

2.0 Attendee List

Karim Alizad	United States Geological Survey
Harris Bienn	The Water Institute of the Gulf
Christine Buckel	National Oceanic and Atmospheric Administration
Kevin Buffington	United States Geological Survey
Charley Cameron	The Water Institute of the Gulf
Juan Cervera	National Oceanic and Atmospheric Administration
Jonathan Clough	Warren-Pinnacle Consulting, Inc.
Renee Collini	Program for Local Adaptation to Climate Effects: Sea-Level Rise
Christopher Esposito	The Water Institute of the Gulf
David Kidwell	National Oceanic and Atmospheric Administration
Connor Levy	National Oceanic and Atmospheric Administration
Sara Martin	Program for Local Adaptation to Climate Effects: Sea-Level Rise
Stephen Medeiros	Embry-Riddle Aeronautical University
James Morris	University of South Carolina
Mary Schoell	National Estuarine Research Reserve Association
Sarah Spiegler	North Carolina State University, North Carolina Sea Grant
Eric White	Coastal Protection and Restoration Authority

3.0 Discussion Summaries

This section includes summaries of the discussions that took place over the course of the workshop, including all pertinent decision points and next steps. Sections 3.1–3.4 summarize discussions from day one, while Sections 3.4–3.7 summarize those from day two.

3.1 Stakeholder Perceptions on Marsh Modeling

On April 11, 2022, marsh modelers convened with the project team to discuss an approach to performing a marsh model retrospective analysis. Before entering these discussions, Sara Martin (PLACE:SLR) and Mary Schoell (National Estuarine Research Reserve [NERR] Association) presented their findings about stakeholder perceptions and use of marsh models. The goals of this session were for marsh modelers to understand 1) how stakeholders use marsh models, 2) what makes a marsh model output most useful for stakeholder applications, and 3) what other model outputs stakeholders would like to see from marsh models.

The presentation contained data obtained through interviews and surveys conducted with stakeholders by Martin, Schoell, and other members of the Project Team. The stakeholders consulted by the project team consisted of researchers and land managers. Stakeholders were interviewed to the point of saturation (i.e., no new trends or information was being gained) for a total of 7 interviews. Following the presentation, the marsh modelers asked questions to further understand how stakeholders use and think about marsh model outputs. Discussion was aided by a posted flipchart page that listed key points from the stakeholder presentation labeled “Managers Want Models That” (Table 2). Following the presentation, marsh modelers were able to ask questions and provide their own perspectives on the use of their models. Feedback from the marsh modelers was captured on a flipchart page labeled “Things for Managers to Know” (Table 3). See the presentation and flipchart page images provided in Appendix A.2 and Appendix D.2-3 respectively.

Table 2. Managers Want Models That...

Managers Want Models That
1. are easy to understand
2. are specific to their managed lands
3. include outputs for various time steps and sea-level rise scenarios
4. are transparent about the input data
5. are clear about what any uncertainty means
6. can analyze management options

Table 3. Things for Managers to Know

Things for Managers to Know
1. Age of National Wetlands Inventory (NWI) Data: NWI vegetation data are outdated in some areas
2. Clarify difference between time steps and (model version) updates
3. Stage of development = generalities: tipping points should be considered by decades, not by year
4. “Elevation capital” = elevation of the marsh above the minimum required for vegetation growth
5. Managers are the tuners for marsh models: their feedback helps modelers to fine tune the models
6. Models are describing vulnerability overall of marsh areas and are not trying to predict the future of exactly what will happen.
7. Models are useful for understanding marsh processes and vulnerabilities

From this discussion, the marsh modelers were able to learn about stakeholder perceptions, identify actions for aiding stakeholders understanding of marsh model outputs, and set goals for continuing to build a dialog between modelers and stakeholders. Furthermore, two *Decision Points* and two *Next Steps* were determined.

3.1.1 Decision Points

1. Stakeholders would be better served by marsh model outputs if they viewed the outputs as a means by which to determine vulnerabilities in a particular marsh, rather than to identify the year that a marsh will reach a “tipping point.”
2. It is important for marsh models to include adequate explanation of source of uncertainty to increase stakeholder confidence in the outputs.

3.1.2 Next Steps

1. Marsh modelers requested a product, such as a two pager, which describes stakeholder perceptions on marsh modeling.
2. The Project Team will also create an extension and outreach product to aid stakeholders understanding of marsh model outputs and provide guidance on applying them based on feedback from the marsh modelers.

3.2 Big Picture Questions for the Marsh Model Retrospective

To prepare the marsh modelers for investigation of the data available for the marsh model retrospective, Trevor Meckley (NOAA) presented details regarding the background of the marsh model retrospective, including why the retrospective is needed, further discussion of the relevance of stakeholder perceptions of marsh models, and summarized the challenges that decision-makers face in applying marsh model outputs. Meckley also highlighted the goals of the workshop and the “big picture questions” that the Project Team seeks to address with the marsh model retrospective. Big picture questions included Example Application Questions and Scoping Questions. See Appendix A.3 and Appendix B.2 for presentation slides and workshop handout listing the questions, respectively.

Following the presentation, the marsh modelers were given time to ask questions and voice any concerns about the marsh model retrospective. The marsh modelers suggested seven considerations for scoping the retrospective process, which were captured on a flipchart (Appendix D.6) and added to the list of scoping questions to consider (Table 4).

As a result of this discussion, the marsh modelers obtained a greater understanding of what the marsh model retrospective will accomplish and the scoping questions for the Project Team and marsh modelers were refined.

Table 4. Considerations for Scoping the Marsh Model Retrospective

Considerations for Scoping the Marsh Model Retrospective	
1.	Need to discern the impact of sea-level rise from other factors (e.g., storms, human changes)
2.	Need to include models in the marsh model retrospective that are transferable to multiple locations
3.	Sites have experienced enough change in sea level to drive change in a marsh
4.	Sometimes things other than sea-level rise, like logging, are driving the marsh changes
5.	Sea-level rise should be a primary driving factor of change in any location selected for inclusion in the marsh model retrospective
6.	The marsh model retrospective analysis will have to avoid sites with punctuated changes or account for any event, management actions, or restorations that have occurred.
7.	River diversions and barrier islands can change the hydrodynamics of distant locations of the system – need to ensure this is acknowledged and addressed if necessary during a retrospective

3.3 Available Data for Performing a Marsh Model Retrospective

Members of the Project Team from The Water Institute of the Gulf guided the marsh modelers through an exploration of the data currently available to run a marsh model retrospective in three locations: Apalachicola NERR, Florida; Grand Bay NERR, Mississippi; and three NERRs in North Carolina (Currituck Banks, Rachel Carson, and Masonboro Island Reserves). The exploration was done through a web-based mapping platform (Figure 1) built by The Water Institute of the Gulf. This mapping tool can be found at <https://arcg.is/0CL9D90>. It included digital elevation model (DEM), tidal datum, land cover, surface elevation table (SET) elevations, and real-time kinematic (RTK) elevation layers for each location across three decades (1990s, 2000s, and 2010s). This data exploration was facilitated by a data inventory handout and a guided exploration activity sheet (see Appendix B.3 and Appendix B.4).




Figure 1. Screenshot of the web mapping platform used for the data exploration by the marsh modelers.

The data exploration was followed by a discussion of the presented data as well as additional data that the marsh modelers identified as beneficial to include. Overall, the marsh modelers found that the tool had an intuitive and easy to use interface and particularly appreciated the inclusion of SET data. The marsh modelers discussed concerns for assessing the results of the marsh model retrospective analysis. Primarily, they discussed how to address the bias of knowing what the retrospective output should look like. For instance, if it is already known that over a twenty-year period there is a 20% decline in vegetation cover how does the project team ensure that the bias does not influence adjustments in the retrospective model?

However, the primary purpose of the data exploration was to determine if there is sufficient data to perform a marsh model retrospective. In addressing that question, the marsh modelers had six concerns (Table 5), identified six data needs (Table 6), and one data want (Table 7). Overall, the group reached two decision points and one next step was identified.

Table 5. Data Concerns

Data Concerns	
1.	Data quality was not clear. Only the most recent data was shown for many of the layers. In particular, the analysis will need historic NWI vegetation data.
2.	NWI data changes over time. The imagery may need to be reclassified so that changes over time reflect actual change rather than changes in classification categories.
3.	SETs are not randomly distributed. Because they are typically located in convenient locations and rarely in "unimpacted" ones, which can give a misleading estimate of rate of change. SET elevations may be biased to eroding edges.
4.	SETs are not routinely monitored. Half of the SETs in North Carolina are not currently monitored due to a lack of funding, so the data may not be available for target time periods. Rather, data from a particular SET may be relevant just to a snapshot in time.
5.	Site selection may have confounding factors. These factors could include salinity, freshwater input, heavy rainfall, etc. This concern could be addressed by including multiple types of sites in the retrospective analysis.
6.	Poor quality or lack of historic DEMs. Performing the marsh model retrospective will require quality DEMs for whatever period is decided on for the analysis. These may not be available, and the need will have to be addressed.

Table 6. Data Needs

Data Needs	
1.	Salinity. The analysis will require long term salinity monitoring.
2.	Total suspended sediments (TSS). A time series of TSS may be needed. This could be acquired from the NERRs System Wide Monitoring Program (SWMP) datasets.
3.	Dated soil cores. These will be required for building a historic DEM.
4.	Water level time series.
5.	Stream gauge data points.
6.	Data trends and points.

Table 7. Data Want

Data Want	
1.	Habitat layer that identifies change. Layers that show drowned forests or ghost trees would be useful.

3.3.1 Decision Points

1. In order to perform the marsh model retrospective, a significant amount of work and funding will be required to support necessary data gathering. In particular, future costs will include processing, adjustment, or standardization of historic NWI data, and gathering the dated soil cores.
2. An objective referee will be needed to evaluate the marsh model retrospective analysis.

3.3.2 Next Steps

1. The marsh modelers requested that the Project Team share the North Carolina and Gulf of Mexico SET inventories with them.

3.4 Day One Wrap-Up

At the close of day one, Renee Collini (PLACE:SLR) reviewed with attendees the accomplishments of the day and provided a brief description of what would be covered in day two of the workshop. The primary outcome from discussions to wrap up day one was a list of outputs to compare from the marsh model retrospective analysis. This list of outputs was determined by the marsh modelers.

Table 8. Marsh Model Retrospective Outputs to Compare

Marsh Model Retrospective Outputs to Compare
Vertical elevations
Horizontal habitat changes
Landscape (holistic output)

3.5 Details for Performing a Marsh Model Retrospective

The second day of the workshop focused on discussion of the details for performing a marsh model retrospective analysis. This included details regarding vegetation input layers, appropriate DEMs, timesteps, sea-level rise, locations, and model uncertainty. The overarching goal of this discussion was to refine the input details for the marsh model retrospective so that the project team and modelers pursue funding to perform the retrospective analysis. Important discussion notes and decision points were captured on flipchart pages throughout the discussion (Appendix D.4-5). Detailed discussion of each topic is provided below.

3.5.1 Vegetation Input Layers

For a model to accurately project changes in marsh habitat, there must first be accurate and reliable layers for existing habitat. However, the project team and marsh modelers identified complications with easily accessible vegetation layers, particularly those available through NWI. Discussions among the project team and the marsh modelers are detailed below.

One of the first concerns raised was that NWI habitat classes have changed over time. Because of this, comparisons of NWI layers over time can be difficult. To remedy this, historic NWI layers will need to be reclassified to ensure that all time periods in the retrospective analysis are comparable. The marsh modelers suggested several ways to complete this task. First habitat re-classification could be accomplished using aerial imagery. To train the reclassification, ground truth data points may be required. However, it is not known if this data exists. Furthermore, historical imagery may be of a lower resolution and therefore the classifications may have to be on the scale of fresh marsh, salt marsh, etc.

In some locations and for some time periods, the historic data for metrics that affect vegetation changes (i.e., accretion, biomass density, TSS, etc.) do not exist. The marsh modelers recommended that site-specific calibration of these metrics could be conducted for the current, existing condition. Then, the relationship between those metrics and vegetation could be assumed to be consistent for historical time periods. By applying those relationship assumptions to historical vegetation layers, metrics such as accretion can be accounted for with reasonable accuracy.

Throughout the discussion, it became clear that each marsh model may have different definitions of “vegetation data.” There was a short pause in the discussion to allow each marsh modeler to define what “vegetation data” means in their model (Table 9). Following that, the project team and marsh modelers discussed what definition should be used for the marsh model retrospective analysis. It was widely agreed, based on what each model already supported for their vegetations layer, that each model could use high and low marsh categories, and possibly tidal-fresh categories for the retrospective analysis. However, the Louisiana ICM is an exception. Because of the low elevation of Louisiana coastal marshes, the elevation-based categories may not apply. However, Eric White (CPRA) suggested that the dominant species could be assigned and be used to create comparable categories for Louisiana marshes.

Table 9. Model Vegetation Data Definitions and Other Important Data

Model	Vegetation Data Definition	Other Important Data
ICM	Uses species level that is then scaled up to the community level	Vertical accretion
Hydro-MEM and MEM	Can be species specific, but generally uses high, mid, and low marsh categories	Biomass and accurate RTK elevations
SLAMM	Needs marsh habitat classes (e.g., high, low, tidal fresh), but can use species mixtures	SET data could be useful, but is not absolutely necessary
WARMER	Uses high and low marsh classifications, but also incorporates dominant species	

The final vegetation topic discussed was how to proceed with the vegetation data layer. Initially, the conversation centered on choosing between NWI or Coastal Change Analysis Program (CCAP) data layers, or an entirely separate dataset. CCAP does not distinguish between high and low marsh, so that would create extra steps in the retrospective process. NWI is likely to provide historical data if provided with a detailed, specific request, but the project team will also need details on the workflow and classification scheme used by NWI. The marsh modelers and project team briefly discussed deriving their own vegetation dataset. However, that would be a more substantial task than building on an existing dataset. It was determined that the project team would begin with historical NWI imagery and reclassify the data to include high, mid, and low marsh while ensuring that each marsh category has a clear definition.

3.5.1.1 Next Steps

- Obtain raw, historical NWI imagery
- Reclassify the imagery to include standardized high and low marsh categories across all time steps; consider other classifications such as mid
- Create clear definitions of high and low marsh, and mid as well if necessary

3.5.2 Digital Elevation Models

An accurate DEM is considered to be a necessity for contemporary marsh models. However, accurate and high resolution DEMs are unlikely to be available for historical time periods due to the lack of LiDAR data, as LiDAR surveys were rarely performed prior to 2007. The marsh modelers first discussed what other historical options are available.

Jonathan Clough (SLAMM) discussed how SLAMM addressed elevation before LiDAR DEMs were available. He suggested that prior approaches would not be applicable for the retrospective because the resulting elevation model would be too coarse to be useful. Even with interpolation, artifacts had big impacts on the marsh model results. Because of the lack of existing historical options, it was decided that the project team and marsh modelers would attempt to build a DEM using historical data. The methodology with which to accomplish this was the focus of conversation for the duration of the DEM discussion. It was acknowledged that building a pre-LiDAR DEM would be a significant undertaking, but the project team and modelers developed a list of the necessary steps and pieces to accomplish the task while also accounting for concerns raised during the discussion.

The first piece that was discussed was the use of dated soil cores to adjust a contemporary DEM to an agreed upon starting point. The marsh modelers agreed that this could be a blunt approach to rolling back time followed by more fine scale modeling. They also agreed that this could be a good approach, but it may be limited by the necessary assumption that the plant community has not changed much over time. To address that limitation, the marsh modelers suggested that historical vegetation data may be able to improve the built DEM resolution. Theoretically, the marsh modelers thought that this would be a good working solution but acknowledged that the accuracy of the resulting DEM may not be sufficient for the marsh model retrospective. Kevin Buffington (WARMER) has experience using dated soil cores to adjust DEMs with success but has not attempted it on a large scale. Jonathan Clough also expressed concern that this may not be valid on dry land adjacent to coastal marshes, which would limit application of the marsh model outputs for marsh migration questions.

Despite the concerns, the marsh modelers agreed that building a new DEM by using soil cores to adjust a good, contemporary DEM would be the best option for the marsh model retrospective analysis.

3.5.2.1 Next Steps

- Choose a good, contemporary DEM
- Obtain data from dated soil cores
- Use soil cores to adjust the DEM to the target time period
- Further increase resolution of the built DEM using historical vegetation data, if needed

3.5.3 Time Steps

To perform a marsh model retrospective analysis, it is critical to agree upon the time steps and time period to consider.

The project team and marsh modelers first discussed what is meant by “time steps.” The marsh modelers pointed out that with modeling there are two types of time steps to consider. There is an internal time step and an external time step. Internal time steps refer to the intervals of data that the model uses in the background processing, like hourly water levels. External time steps refer to the time steps presented in the model outputs. The project team clarified that for the purposes of this discussion, the marsh modelers should consider external time steps.

To begin selection of time steps to use in the marsh model retrospective, the discussion referred to what stakeholders could be expected to want to see in model outputs. Because stakeholders would be most interested in intermediate time spans for the comparison, it was agreed that five-year time steps would be ideal. However, the marsh modelers felt that this would primarily depend on data availability. Point elevations, rather than landscape ones (i.e., LiDAR DEMs) would be most useful for achieving five-year time steps. Overall, the project team and marsh modelers agreed that five-year time steps would be possible. However, a starting and ending time point was not agreed upon. Rather, it was agreed that data availability would have to be further explored at target locations to make this decision.

3.5.3.1 Next Steps

- Obtain historical point elevations.
- For the marsh model retrospective analysis, the outputs will have five-year time steps.

3.5.4 Sea-Level Rise

Sea-level rise data are necessary in considering changes to coastal marshes in marsh models. While the retrospective analysis will use known inputs, sea-level rise is recorded in linear and non-linear trends. To determine which type of sea-level rise data to use in the marsh model retrospective, the marsh modelers discussed a few points.

First, Renee Collini presented information about the difference between the linear and non-linear sea-level rise trends. Based on her presentation, the marsh modelers needed clarification on using the observed data versus using a sea-level rise trend. If we have known data, then that would seem ideal for model applications, especially considering the high amount of variability in water levels over time. However, because the retrospective is intended to run from a starting point and forecast unknown conditions, a sea-level rise trend would be most appropriate.

Next, the marsh modelers discussed the choice between the linear and non-linear sea-level rise trends. They determined that it would be most useful to choose the trend to use based on the nearest tide station to the targeted locations. Each model would perform the retrospective based on the curve that has the best fit at each location. However, the marsh modelers also pointed out that this exercise could also be useful for identifying areas of uncertainty in each model if compared to model runs using the actual observed data. Therefore, outside of the marsh model retrospective analysis, the marsh modelers will run their model with observed data to support extension and outreach to stakeholders.

3.5.4.1 Next Steps

- Identify the sea-level curve with the best fit (linear or nonlinear) at tide stations nearest the target locations.
- Use the selected curve to run the models in the marsh model retrospective analysis.
- To create extension materials for stakeholders, compare the results of the analysis with another model run using observed sea-level rise data.

3.5.5 Locations

Throughout much of the discussion during day two of the workshop, it was mentioned that data availability by location may be a limitation for different input parameters. For this reason, the discussion moved towards narrowing down location choices for the marsh model retrospective analysis. This process began with the marsh modelers suggesting locations that they deemed to fit two criteria: 1) the location has a wealth of data, and 2) there has been enough change in the marsh that the models would be able

to describe the change. Each location was then discussed to determine what data may be easily accessible.

3.5.5.1 Grand Bay NERR

The Grand Bay NERR was suggested because it has been well studied, several of the models represented in at the workshop have already been run there, and the marsh modelers believe that data would be easy to obtain. Grand Bay would also serve as a good representative of how the models perform in a microtidal system.

3.5.5.2 Apalachicola NERR

Apalachicola NERR, like Grand Bay NERR, has been well studied and several of the marsh models have already been ran there. However, Apalachicola is a relatively complex system that could create difficulties in the marsh model retrospective analysis. There have been many changes in the Apalachicola NERR that may have a variety of drivers. This would make it difficult to distinguish the effects of sea-level rise from other drivers.

3.5.5.3 North Carolina Reserve System

The North Carolina Reserve sites have been well studied (e.g.vegetation surveys, SETS, water level data) and were included in the data mining that was explored in day one of the workshop, so it is known that there is a lot of existing data in the region. The marsh modelers also pointed out that these locations would serve as a good comparison to the microtidal locations along the northern Gulf of Mexico that had already been suggested. However, some marsh models have not been run in this area.

3.5.5.4 Louisiana

Coastal marshes in Louisiana have experienced considerable amounts of change, and there is a breadth of data available. In particular, the marsh modelers specified available vegetation surveys, SETs, accretion, and water level data. USGS also has regularly updated aerial imagery that may be useful for reclassifying vegetation data layers for the marsh model retrospective analysis. Eric White indicated that CPRA is interested in doing a rigorous hindcast in the region, which could be leveraged for this effort. There were two concerns raised by the marsh modelers. First, the complexity of Louisiana marshes could prove difficult. Second, a specific location within Louisiana was not agreed upon during the workshop.

3.5.5.5 Plum Island, MA

Plum Island is a marsh that is data rich. In particular, James Morris and Karim Alizad expressed high levels of familiarity with the data available in that area. They also indicated that there has already been a lot of change in marsh extent at Plum Island. Plum Island could also be a representative location for marshes along the east coast of the United States.

3.5.5.6 Sacramento Delta

The marsh modelers were asked to suggest locations on the west coast in addition to those already suggested on the east and Gulf coasts of the United States. The Sacramento Delta was suggested as the best-known choice. However, areas of significant change may be hard to find in the area. Elkhorn Slough was suggested, but otters have played a key role in driving the change there. While there are data rich areas in the region, selecting one with significant change may be a challenge.

After discussion of the possible locations, the marsh modelers were asked to indicate their preference for where to run the marsh model retrospective by placing sticky dots next to their top three locations on the flipcharts. When they completed that task, the votes were tallied to decide which three locations to focus on for this effort. Through this, it was decided that the project team and marsh modelers would move

forward in Louisiana, Grand Bay NERR, and Plum Island, MA. Primarily, these locations were selected due to the availability of data and the breadth of complexities and variables that the locations cover. The voting tally can be seen in Table 10 below or on the flip chart images (Appendix D.5).

Table 10. Votes for Marsh Model Retrospective Locations

Location	Votes
Apalachicola NERR	3
Grand Bay NERR	5
Louisiana	4
North Carolina Reserves	1
Plum Island, MA	5

3.5.5.7 Next Steps

- The marsh modelers decided to focus on Louisiana, Grand Bay NERR, and Plum Island, MA for the marsh model retrospective.
- A specific location will need to be selected in Louisiana.
- The project team and marsh modelers will work to inventory the available data at each of the three locations to ensure their applicability in the marsh model retrospective.

3.5.6 Model Uncertainty

Lastly, the marsh modelers discussed how to address model uncertainty based on stakeholder feedback. Stakeholders indicated that sources and meaning of uncertainty should be clear so that they can understand how to account for it in their decision making (for more information, see the Stakeholder Perceptions Presentation in Appendix A.2). The marsh modelers first defined the uncertainty in their models (Table 11). Then, they decided that for the marsh model retrospective they will set boundaries on the uncertainty based on the various models and the range of conditions that the models will use. The marsh modelers also pointed out that this effort for the marsh model retrospective could be used as an opportunity to characterize the uncertainty for stakeholders.

Table 11. Sources of Uncertainty in Models

Model	Sources of Uncertainty
SLAMM	This model has some uncertainty bounds around inputs (including DEMs) that help to generalize the confidence intervals and relative vulnerability on marsh presence likelihood.
WARMER	This model samples from a distribution of accretion rates, decomposition, and other factors and runs Montecarlo simulations to get a confidence level.
ICM	This model has more scenario-based uncertainty. It has some model validation to generate statistics and then perturbs the model.
Hydro-MEM	No details about model uncertainty are provided in model outputs.
MEM	This model uses the Montecarlo approach with uncertainty

3.5.6.1 Next Steps

- Once the data are defined, bounds of uncertainty and the range of conditions that will be used in the marsh model retrospective analysis will be set.
- This effort will be leveraged to create extension products for stakeholders to explain uncertainty in model outputs.

3.6 Discussion Summary: Revisiting the Marsh Model Retrospective Framework

Following the discussions about what details to include in the marsh model retrospective analysis, the modelers and project team returned to the questions regarding framework to ensure that all the decisions that had been made will support the example application and scoping questions listed at the beginning of the workshop (questions are included in Section 3.2, and in the handout provided in Appendix B.2).

The group first reviewed the scoping questions. Questions for which there was discussion are listed below, with specific discussion points noted.

1. What site specific variables do we need to evaluate for their influence on model performance?
 - a. Estuary type and characteristics
 - i. The modelers stated that the locations selected cover both a range of estuary types and site complexities that will sufficiently test the capabilities of the models.
2. How do we evaluate marsh model performance? What outputs do we compare?
 - a. The modelers detailed two ways that the outputs could be compared: cell-by-cell in the output maps, by percent area, or both.
 - b. Cell-by-cell is likely to be more reflective of the starting conditions and cell sizes.
 - c. Percent land cover change may be more informative, but the modelers were unsure if it would answer questions related to where a model performs well.
 - i. Instead of cell-by-cell analysis, the outputs could be divided into sub-domains, in which model “hits or misses” could be further explored.
 - d. For the purposes of comparison, areas where the model outputs are predicting similar vulnerabilities should be highlighted.
 - e. The comparison can be depicted using a histogram of vegetation types or classes.
3. Do we have enough data, and the needed data, to perform the retrospective analysis and to answer the research questions right now?
 - a. The modelers agreed that there is too much uncertainty on the data availability to answer this question.

The group then reviewed the example application questions. The modelers were asked if the questions could be answered in the three selected locations based on what had been scoped throughout the workshop. The modelers primarily discussed the use of the retrospective in addressing questions of marsh migration. They expressed concerns that the limitations of a constructed DEM, as discussed in Section 3.5.2, would undersell migration potential and model certainty. This would need to be addressed in communications with the stakeholders about the retrospective analysis.

3.7 An Approach and Timeline for the Marsh Model Retrospective

The final discussion of the workshop was focused on gathering the information gained from the two-day workshop to outline an approach for performing the marsh model retrospective. This included listing next steps, assigning who will be responsible for each task, estimating costs, and setting a timeline for completing the tasks. The decisions were captured on a flipchart (Appendix D.7) and are outlined below. The purpose for each task was discussed in detail throughout the workshop and can be found in Section 3.5.

Task:	Acquire NWI Data
Responsibilities:	Christine Buckel and Trevor Meckley will reach out to the United States Fish and Wildlife Service to request NWI data layers.
Timeline:	Approximately two months
Estimated Costs:	No associated costs

Task:	Reclassify NWI Data
Responsibilities:	Renee Collini will facilitate discussions about classifications for the data. The project team will ask for volunteers to participate in the discussions once the NWI data is in hand. The Water Institute of the Gulf expressed interest in doing the data classification once funding is available.
Timeline:	6–8 months for the entire process once the data is in hand
Estimated Costs:	Funding will be required to pay for someone's time to do the classification of the NWI data.

Task:	Gather DEM data for the selected locations
Responsibilities:	Responsibilities were discussed by site: Louisiana: Eric White will lead the effort for this site. Plum Island: Karim Alizad believes that Matt Kirwan has the existing DEM for Plum Island and will reach out to him. Grand Bay: The project team already has access to the DEM for Grand Bay.
Timeline:	2–4 months for all three sites,
Estimated Costs:	There are no estimated costs associated with this task.

Task:	Pick a specific location for Louisiana.
Responsibilities:	Eric White and Christopher Esposito will lead the selection of a site in Louisiana
Timeline:	Approximately 2 months because some data will need to be found, but not analyzed, before final site selection occurs.
Estimated Costs:	No associated costs.

Task:	Identify existing soil cores.
Responsibilities:	Trevor Meckley, James Morris, and Eric White will work together to first define what soil core data is needed for the retrospective analysis The leads for identifying soil cores differed based on location.

Louisiana: Eric White
Plum Island: Karim Alizad and James Morris
Grand Bay: Karim Alizad

Christine Buckel will be the point of contact for assuring this task is completed on time.

Timeline: Approximately 4 months.

Estimated Costs: No associated costs.

Task: Acquire Sea-Level Rise Data.

Responsibilities: Renee Collini will lead this effort.

Timeline: Renee indicated that this task can be done quickly once the final site selection is completed.

Estimated Costs: No associated costs.

Task: Acquire RTK data.

Responsibilities: The lead for this task differs by location:

Louisiana: Eric White
Plum Island: Karim Alizad
Grand Bay: Renee Collini

Timeline: 8 months

Estimated Costs: No associated costs.

Task: Build the historical DEMs

Responsibilities: Due to the complex nature of this task, the project team and modelers believe that someone will have to be hired to complete it.

Timeline: 18 months

Estimated Costs: Money will be required to hire someone to complete this task.

Task: Obtain new soil cores to fill in any gaps in the existing soil cores.

Responsibilities: Due to the complex nature of this task, the project team and modelers believe that someone will have to be hired to complete it.

Timeline: 2–5 years for full collection and analysis.

Estimated Costs: Money will be required to hire someone to complete this task.

Task: Determine specific model costs

Responsibilities: Trevor Meckley will reach out to each modeler to determine their specific model costs.

Timeline: 1 month.

Estimated Costs: No associated costs.

3.7.1 Next Steps

Following the discussion of the timeline and specific tasks to be completed before a marsh model retrospective analysis could be conducted, Renee Collini led discussion of the next steps to follow the workshop.

To ensure continued communication and that updates are shared with the entire project team, emails will be sent out quarterly. Virtual meetings will be held semi-annually or as needed based on project progress. These meetings will be organized using online polling to select dates, and available data will be sent to the project team ahead of any meetings. This will ensure that valuable time is spent on discussion in the meetings rather than extensive data presentations.

David Kidwell expressed concerns that the estimated timeline for the retrospective analysis (5 years) is too long; the need for the retrospective is too pressing. To hasten the timeline, the project team and modelers agreed to address the following tasks within six months of the workshop completion:

- Assess availability of existing soil cores
- Identify existing RTK data
- Obtain existing NWI data
- Obtain existing DEM data
- Pick a specific Louisiana location
- Determine costs for running each of the models

This list will form the checklist of tasks to be completed before the first virtual meeting.

4.0 Appendices

Appendix A: Workshop Presentations

A1: Introduction to the Workshop

Marsh Model Retrospective Workshop

Beaufort Hotel | April 11-12, 2022 (finally)

Packets

- ▶ Agenda
- ▶ Meeting Logistics
- ▶ Question List
- ▶ Data Inventory Sheet
- ▶ Activity Sheet

Agenda - Day One

- ▶ Background & Setting the Stage
 - ▶ Overview - Goals and Objectives
 - ▶ Stakeholder Input
 - ▶ What is a Retrospective
- ▶ Scoping Question
- ▶ Break
- ▶ Review Data Collected So Far
 - ▶ Introduction
 - ▶ Data Exploration
- ▶ Introduce Technical Details
- ▶ Review Scoping Questions
- ▶ **Dinner - Moonrakers**

Agenda - Day Two


- ▶ Review Day One
 - ▶ Progress - what we found
 - ▶ Reflection
- ▶ Technical Details
 - ▶ Veg Input Data
 - ▶ DEMs
 - ▶ Time Steps
- ▶ Break
- ▶ Technical Details
 - ▶ SLR
- ▶ Locations
 - ▶ Anything else?
- ▶ Revisit Scoping Questions
- ▶ Revisit Example Application Questions
- ▶ Details of the Retrospective
 - ▶ Working **lunch** if need-be
- ▶ Next Steps & Wrap-Up
- ▶ **Field Trip**

A2: Stakeholder Perceptions of Marsh Models



In 2018, marsh modelers and stakeholders met to compare and discuss the outputs of four marsh habitat change models.

Marsh Model Retrospective



Why do stakeholder perspectives matter?

Stakeholders are the end-users



Gathering Stakeholder Perspectives



The stakeholder interviews were driven by three main questions:

- How do stakeholders use marsh model outputs?
- If they are not using model outputs, then why not?
- What makes a marsh model output more applicable to stakeholders?

Interviews were conversational and generally covered 9 questions

1. How do you use marsh models in your work?
2. What models do you use most often?
3. What environmental inputs are important for models to consider?
4. How do you resolve differences between model outputs?
5. What time steps are most useful for your decision making?
6. What level of parcel averaging are you comfortable with?
7. What model outputs would be most useful other than marsh presence/absence?
8. How likely are you to trust the results of the retrospective analysis?
9. How should we report the results of the retrospective?

Interviews were conversational and generally covered 9 questions


1. How do you use marsh models in your work?
2. What models do you use most often?
3. What environmental inputs are important for models to consider?
4. How do you resolve differences between model outputs?
5. What time steps are most useful for your decision making?
6. What level of parcel averaging are you comfortable with?
7. What model outputs would be most useful other than marsh presence/absence?
8. How likely are you to trust the results of the retrospective analysis?
9. How should we report the results of the retrospective?

Interviews were conversational and generally covered 9 questions


1. How do you use marsh models in your work?
2. What models do you use most often?
3. What environmental inputs are important for models to consider?
4. How do you resolve differences between model outputs?
5. What time steps are most useful for your decision making?
6. What level of parcel averaging are you comfortable with?
7. What model outputs would be most useful other than marsh presence/absence?
8. How likely are you to trust the results of the retrospective analysis?
9. How should we report the results of the retrospective?

Results


How do stakeholders use marsh models?



How do stakeholders use marsh models?




How do stakeholders use marsh models?






Why aren't stakeholders using marsh models?

- Does not apply to their work
- Not aware of available outputs
- Do not have the capacity to use them
- Do not need marsh model outputs... yet!
- Does not trust model outputs




What works well in marsh model outputs?


- Easy to understand
- Is a good scale for application
- Have a variety of SLR scenarios
- Wetland migration outputs
- Considers many input parameters
- Shows change over time



What time steps are most useful to stakeholders?




What time steps are most useful to stakeholders?




What kinds of outputs, other than marsh presence/absence, would most useful to stakeholders?

- Community level changes
- Ecosystem shifts
- Information about inputs
- Ecosystem function
- Marsh age
- Elevation
- Invasive species
- High water
- Historic shoreline position
- Uncertainty




What kinds of outputs, other than marsh presence/absence, would most useful to stakeholders?


- Community level changes
- Ecosystem shifts
- Information about inputs
- Ecosystem function
- Marsh age
- Elevation
- Invasive species
- High water
- Historic shoreline position
- Uncertainty




How likely are stakeholders to trust the results of the retrospecti



How likely are stakeholders to trust the results of the retrospecti



How likely are stakeholders to trust the results of the retrospective



Key takeaways from the conversations with stakeholders

Key takeaways from the conversations with stakeholders

While most stakeholders do use models in their decision making, there is a segment that do not.

Key takeaways from the conversations with stakeholders

The more outputs and scenarios a model considers increases its useability for stakeholders, especially if it can be applied cheaply and easily for their managed lands.

Key takeaways from the conversations with stakeholders

Many stakeholders expressed concerns about transparency in model processes and surrounding uncertainty in the outputs.

How does this apply to this workshop?


Example Application Questions

1. What model should I use for my marsh type?
2. What is the utility of different model output types for different management decisions?
3. How certain are the predictions and are there areas where we are more confident?



Questions?

A3: Background on the Marsh Model Retrospective




 **NCCOS** NATIONAL CENTERS FOR
COASTAL OCEAN SCIENCE

Background on Marsh Model Retrospective Analysis

SCIENCE SERVING COASTAL COMMUNITIES

Retrospective analysis uses historical data to predict known conditions



Mogensen and Rogers, 2018 | DOI:10.1038/s41598-018-19695-2

Marsh manager/planners have questions on application



- Where should we acquire land or protect land (policy) or how should realign coastal infrastructure to ensure marshes exist in the future?
- How will management actions affect my marsh?
 - Controlled burn
 - Fronting structure to protect against erosion
 - Changing land elevation through placement of sediment
 - Estuary mouth opening (or allow to remain closed)
- These application questions have led to questions on what model should I use for my marsh and how accurate (certain) are the results?



We hope to advance the entire field with a retrospective analysis

- **Reduce end user confusion in model outputs**
- **Determine where and when models perform well to guide advancement**
- **Inform future investment**
- **Build a library of marsh data and outputs**
- **Build a modeler community of practice**

Questions

A4: Introduction to Data Exploration





North Carolina

Sites

North Carolina sites include the Currituck Banks, Rachel Carson, and Masonboro Island NERRs.

Elevation

2010s representative DEMs for all locations were sourced from the NOAA NCEI published between 2018 and 2019 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE.

2000s representative DEMs for all locations were sourced from the USGS NED published in 2003 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. The Coastal Relief Model (CRM) was used as the bathymetry resource for both the Rachel Carson and Currituck Banks sites. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.

1990s representative DEMs for all locations were sourced from the NOAA NCEI CRM Volume 2 published in 1998 at a resolution of 3 arc seconds (approximated to 90 meters). Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW) with source topography in NAVD88.

Tidal Datum Gauges

- Currituck Banks - 8651370, Duck, NC
- Rachel Carson - 8656483, Beaufort, Duke Marine Lab, NC
- Masonboro Island - 8658163, Wrightsville Beach, NC



[Navigate Here](#)

North Carolina

Sites

North Carolina sites include the Currituck Banks, Rachel Carson, and Masonboro Island NERRs.

Elevation

2010s representative DEMs for all locations were sourced from the NOAA NCEI published between 2018 and 2019 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE.

2000s representative DEMs for all locations were sourced from the USGS NED published in 2003 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. The Coastal Relief Model (CRM) was used as the bathymetry resource for both the Rachel Carson and Currituck Banks sites. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.



1990s representative DEMs for all locations were sourced from the NOAA NCEI CRM Volume 2 published in 1998 at a resolution of 3 arc seconds (approximated to 90 meters). Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW) with source topography in NAVD88.


Tidal Datum Gauges

- Currituck Banks - 8651370, Duck, NC
- Rachel Carson - 8656483, Beaufort, Duke Marine Lab, NC
- Masonboro Island - 8658163, Wrightsville Beach, NC



[Navigate Here](#)





SOUTHEAST
Climate Adaptation Science Center

Stakeholder Data Investigation Tool

This web mapping application was developed to inform a retrospective marsh modeling workshop occurring in Spring 2022. It leverages a combination of digital elevation models (DEMs), digitized tidal datums, land cover data, and point locations for SET and RTK resources.

Data is presented for several NERR system locations in Mississippi, Florida, and North Carolina. Pages and data resources are represented in decadal bins split by state with source data assigned to a representative decade.

North Carolina

Sites
North Carolina sites include the Currituck Banks, Rachel Carson, and Masonboro Island NERRs.

Elevation
2010s representative DEMs for all locations were sourced from the NOAA NCEI published between 2018 and 2019 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE.

2000s representative DEMs for all locations were sourced from the USGS NED published in 2003 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. The Coastal Relief Model (CRM) was used as the bathymetry resource for both the Rachel Carson and Currituck Banks sites. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.

1990s representative DEMs for all locations were sourced from the NOAA NCEI CRM Volume 2 published in 1998 at a resolution of 3 arc seconds (approximated to 90 meters). Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW) with source topography in NAVD88.

Tidal Datum Gauges



- Currituck Banks - 8651370, Duck, NC
- Rachel Carson - 8656483, Beaufort, Duke Marine Lab, NC
- Masonboro Island - 8658163, Wrightsville Beach, NC



Florida



Sites
Florida sites are limited to the Apalachicola NERR.


Elevation
2010s representative DEMs were sourced from the NOAA NCEI published in 2019 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE.


2000s representative DEMs were sourced from the USGS NED published between 2007 and 2010 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Only topographic data is available for this location and decade.



































Stakeholder Data Investigation Tool

North Carolina

2000s

North Carolina NERR Boundaries

NAME

- Currituck Banks
- Masonboro Island
- Rachel Carson

North Carolina RTK

North Carolina SET

Elevation Trend

- Positive
- Negative
- No Trend

Select a Different Decade

'90s '00s '10s

SECASC Data Resource Reporting - North Carolina

Use the map annotation and comments dialogue on this page to provide the location and details of any additional information that you think might be pertinent to this project.

Please submit one record at a time. After submitting a response this page will reload to provide you with the opportunity to submit another record.

Beaufort, NC, USA

Perimeter: 212.89 Feet Area: 0.08 Acres

Comments

Please provide us with additional information regarding your response.

Huge Japanese Maple in the courtyard of a local dive. |

946

Submit

View a Different Location

Alabama & Mississippi

North Carolina

Florida

Stakeholder Data Investigation Tool

North Carolina

2000s

North Carolina NERR Boundaries

NAME

- Currituck Banks
- Masonboro Island
- Rachel Carson

North Carolina RTK

North Carolina SET

Elevation Trend

- Positive
- Negative
- No Trend

Select a Different Decade

'90s '00s '10s

SECASC Data Resource Reporting - North Carolina

Thank you for taking the time to submit a detailed response.

This page will reload momentarily if you would like to make another submission.

View a Different Location

Alabama & Mississippi

North Carolina

Florida

Stakeholder Data Investigation Tool

North Carolina

2000s

SEASC Data Collection - North Carolina - Stakeholder Responses

Age, in hours, from time created to 4/14/2022, 12:00 PM

- > 313.16
- 313.108
- < 313.052

North Carolina NERR Boundaries

NAME

- Currituck Banks
- Masonboro Island
- Rachel Carson

North Carolina RTK

North Carolina SET

Elevation Trend

- Positive
- Negative

Select a Different Decade

'90s '00s '10s

Layers

Layer Legend

- SEASC Data Collection - North Carolina - Stakeholder
- Hide layer
- North Carolina NERR Boundaries
- North Carolina RTK
- North Carolina SET
- 2000s Representative Tidal Contours
- 2000s Representative DEM (waters NAVD83, NED)
- National Wetlands Inventory - North Carolina
- National Wetlands Inventory - North Carolina - Metadata
- Estuarine CCAP Landcover - North Carolina

Zoom to

Survey Response

Comments

Huge Japanese Maple in the courtyard of a local dwe.

Last edited by harris.benn on 4/11/2022, 9:31 AM.

SEASC Data Resource Reporting - North Carolina

Use the map annotation and comments dialogue on this page to provide the location and details of any additional information that you think might be pertinent to this project.

Please submit one record at a time. After submitting a response this page will reload to provide you with the opportunity to submit another record.

Find address or place

Powered by Esri

Comments

Please provide us with additional information regarding your response.

Submit

View a Different Location

Alabama & Mississippi

North Carolina

Florida

Stakeholder Data Investigation Tool

North Carolina

2000s

North Carolina NERR Boundaries

NAME

- Currituck Banks
- Masonboro Island
- Rachel Carson

North Carolina RTK

North Carolina SET

Elevation Trend

- Positive
- Negative
- No Trend

2000s Representative Tidal Contours

Currituck Banks, Rachel Carson, Masonboro Island

Tidal Datum

- MLLW
- MLW
- MSL
- MHW
- MSL+0.5

Select a Different Decade

'90s '00s '10s

100 m

300 ft

QUESTIONS?


View a Different Location

Alabama & Mississippi

North Carolina


Florida

A5: Marsh Model Retrospective Framework



NOAA's COMT as an example

- NOAA's Coastal and Ocean Modeling Testbed (COMT) offers a general framework for how we could collaborate
- COMT creates a dataset for evaluating models to better understand them for operational running of models
- We may even be able to use their platform.



<https://ioos.noaa.gov/project/comt/>

Our testbed will focus on improving application

Our Goal is to run a test bed that fosters organized model advancement and sharing and **does not crown an “operationally” superior model.**

Marsh modeling is not ready for operations and will never involve model runs as regularly as the models COMT has considered.

We expect modelers to all be involved in running models and hope to design the retrospective analysis in a way to benefit modelers.

We are going to ask for input across two areas

- **Scoping questions** to inform the approach to accomplishing the analysis
- **Technical details** on data needs that will inform how we scope the analysis.

Known Conditions



Model 1



Model 2



Model 3



Questions for scoping the retrospective analysis:

- What site specific variables do we need to evaluate for their influence on model performance?
 - Estuary type and characteristics to consider?
 - Data quality of data inputs to consider?
- How do we evaluate marsh model performance?
 - What outputs do we compare?
- Do we have enough data to run a retrospective right now?
- Are there other scoping aspects to consider?

Appendix B: Handouts

B1: Participant Agenda

Marsh Model Retrospective Workshop Day 1

April 11-12, 2022 | The Beaufort Hotel | Beaufort, North Carolina

12:00 pm	Lunch Grab some lunch provided by 34° North
1:00 pm	Welcome and Introductions Get to know everyone
1:20 pm	Introduction to the Day Establish ground rules, go over the agenda for the day, and review the reasons for performing a marsh model retrospective analysis
1:45 pm	Stakeholder Perceptions on Marsh Models Learn why stakeholders use marsh model outputs
2:05 pm	Big Picture Questions for the Marsh Model Retrospective Review the questions that the retrospective will seek to answer
2:20 pm	Break With snacks!
2:35 pm	The Work So Far Review the data that has been accumulated so far
2:50 pm	Data Exploration and Discussion Explore the data using this link: https://arcg.is/0CL9D90
3:45 pm	The Marsh Model Retrospective Framework Revisit the Big Questions again
4:15 pm	Wrap-up Review the accomplishments of the day and prepare for Day 2
4:25 pm	Adjourn

5:30 pm

Dinner at Moonrakers

Marsh Model Retrospective Workshop Day 2

April 11-12, 2022 | The Beaufort Hotel | Beaufort, North Carolina

9:00 am

Welcome & Introduction to Day 2

Review the agenda for Day 2 of the workshop and get everyone on the same page before diving into the details

9:35 am

Details of the Marsh Model Retrospective (Part 1)

Dig into the details of when, where, and how to run the analysis

10:15 am

Break

With snacks!

10:30 am

Details of the Marsh Model Retrospective (Part 2)

More discussion of details for the retrospective analysis

11:15 am

Return to the Big Picture Questions

Now that we discussed the details, how do you feel about the Big Questions?

11:30 am

Putting All of the Pieces Together

Determine what the Marsh Model Retrospective will look like

12:00 pm

Working Lunch

Next steps for the Marsh Model Retrospective

12:50 pm

Workshop Wrap-Up

Recap and review what has been accomplished and next steps

1:00 pm

End of Workshop

Thank you for your input and hard work!

2:00 pm

Field Trip to the Rachel Carson Reserve

Meet at the docks across from Beaufort Hotel for a tour by the reserve staff

B2: Questions for Marsh Model Retrospective

Questions for the Marsh Model Retrospective

Example Application Questions – These are examples of questions that we hope to answer by performing a marsh model retrospective. These questions will inform the framing of much of the discussion throughout the workshop.

1. What model should I use for my marsh type?
2. What is the utility of different model output types for different types of management decisions?
3. How certain are the predictions and are there areas we are more confident?

Scoping Questions – These questions will inform the approach to accomplishing the analysis.

1. What site specific variables do we need to evaluate for their influence on model performance?
 - a. Estuary type and characteristics
 - b. Quality of data inputs
2. How do we evaluate marsh model performance? What outputs do we compare?
3. Do we have enough data and the right data to perform the retrospective analysis and to answer the research questions right now?
4. Are there any other scoping questions that we need to consider?

B3: Data Inventory

Data Inventory NORTH CAROLINA

DEMS

Note: Rows shaded in orange indicate no bathymetry, yellow shading indicates that the vertical datum was something other than NAVD 88 for bathymetry.

Masonboro Island

Source	Resolution	Tidal Datum	Relative Tidal Datum	Vertical Datum	Vertical Units	Date Published	Decade Represented	Notes
NOAA NCEI	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2019	2010s	Topobathy. Site mosaicked from 2 individual 15 minute x 15 minute rasters. Bathymetric and topographic data utilized for DEM creation originate from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE. DEMs are referenced vertically to NAVD 88 with vertical units in meters.
USGS NED	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2003	2000s	Topography only. Site mosaicked from 2 individual rasters. The NED is derived from diverse source data that are processed to a common coordinate system and unit of vertical measure.
NOAA NCEI CRM Vol. 2 (THREDDs)	~90m - 3 arc second MLLW (EPSG: 5866)	MLLW (EPSG: 5866)	MSL	NAVD88 for topography	Meters	1998	1990s	Topobathy. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW). Source topographic data were in NAVD 88. The differences between these datums are less than the vertical accuracy of the CRM, so you can assign MSL to the CRM if you like, just recognize that the elevation values may not be as accurate as you might like or need. Assume a vertical accuracy no better than 1 meter for any elevation values in the CRM.

Rachel Carson Reserve

Source	Resolution	Tidal Datum	Relative Tidal Datum	Vertical Datum	Vertical Units	Date Published	Decade Represented	Notes
NOAA NCEI	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2019	2010s	Topobathy. Site mosaicked from 2 individual 15 minute x 15 minute rasters. Bathymetric and topographic data utilized for DEM creation originate from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE. DEMs are referenced vertically to NAVD 88 with vertical units in meters.
USGS NED	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2003	2000s	Topobathy, bathymetry from CRM. Site mosaicked from 2 individual 15 minute x 15 minute rasters. The NED is derived from diverse source data that are processed to a common coordinate system and unit of vertical measure. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.
NOAA NCEI CRM Vol. 2 (THREDDs)	~90m - 3 arc second MLLW (EPSG: 5866)	MLLW (EPSG: 5866)	MSL	NAVD88 for topography	Meters	1998	1990s	Topobathy. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW). Source topographic data were in NAVD 88. The differences between these datums are less than the vertical accuracy of the CRM, so you can assign MSL to the CRM if you like, just recognize that the elevation values may not be as accurate as you might like or need. Assume a vertical accuracy no better than 1 meter for any elevation values in the CRM.

Currituck Banks

Source	Resolution	Tidal Datum	Relative Tidal Datum	Vertical Datum	Vertical Units	Date Published	Decade Represented	Notes
NOAA NCEI	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2019	2010s	Topobathy. Site mosaicked from 2 individual 15 minute x 15 minute rasters. Bathymetric and topographic data utilized for DEM creation originate from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE. DEMs are referenced vertically to NAVD 88 with vertical units in meters.
USGS NED	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2003	2000s	Topobathy, bathymetry from CRM. Site mosaicked from 2 individual 15 minute x 15 minute rasters. The NED is derived from diverse source data that are processed to a common coordinate system and unit of vertical measure. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.
NOAA NCEI CRM Vol. 2 (THREDDS)	~90m - 3 arc second MLLW (EPSG: 5866)	MLLW (EPSG: 5866)	MSL	NAVD88 for topography	Meters	1998	1990s	Topobathy. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW). Source topographic data were in NAVD 88. The differences between these datums are less than the vertical accuracy of the CRM, so you can assign MSL to the CRM if you like, just recognize that the elevation values may not be as accurate as you might like or need. Assume a vertical accuracy no better than 1 meter for any elevation values in the CRM.

DEM CORRECTIONS

Data for RTK surveys conducted by Brandon Phuckett available for the following locations and dates:

Location	Date
Carrot Island	09/2018-08/2019
Beaufort (site: middle marsh)	2008, 2012, 2015, 2017
Wilmington (site: Zeke's Island)	2013, 2016, 2017, 2019, 2021

ECOLOGICAL DATA

Ecological Data is also available from the National Wetlands Inventory

Location	Type	Date	Source
Beaufort, NC	Point Salinity Dominant Plan Communities and Vegetation Community Types	2015-2020	NOAA / NCCOS-Beaufort
Theodore Roosevelt Natural Area	Point Salinity Dominant Plan Communities and Vegetation Community Types	2016	UNC / Voss
Cedar Island NWR	Point Salinity Dominant Plan Communities and Vegetation Community Types	2016	UNC / Voss

SEA LEVEL TRENDS


Nonlinear trends for Beaufort, NC

Created using data from the Interagency Sea Level Rise Scenario Tool:

https://sealevel.nasa.gov/task-force-scenario-tool?psmsl_id=2295


Technical Report available:

<https://oceanservice.noaa.gov/hazards/sealevelrise/sealevelrise-tech-report-sections.html>



Relative Sea Level (linear) trends for Beaufort, NC

From NOAA: https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8656483



GAUGE DATA

Currituck Banks

Present Installation Start: 27-Jul-88	Datum	Elevations on NAVD88	Elevations on Mean Lower Low Water	Elevations on Mean Sea Level
Station Home Page: 8651370, Duck, NC	MHHW	0.457	1.124	0.585
Station Datum Definitions: 8651370, Duck, NC	MHW	0.36	1.027	0.488
	MSL	-0.128	0.539	0
RSLR: 8651370, Duck, NC	MLW	-0.623	0.044	-0.496
	MLLW	-0.667	0	-0.539

Rachel Carson Reserve

Present Installation Start: 27-Jul-88	Datum	Elevations on NAVD88	Elevations on Mean Lower Low Water	Elevations on Mean Sea Level
Station Home Page: 8656483, Beaufort, Duke Marine Lab, NC	MHHW	0.455	1.078	0.557
Station Datum Definitions: 8656483, Beaufort, Duke Marine Lab, NC	MHW	0.358	1.991	0.47
	MSL	-0.122	0.521	0
RSLR: 8656483, Beaufort, Duke Marine Lab, NC	MLW	-0.59	0.043	-0.478
	MLLW	-0.633	0	-0.521

Masonboro Island

Present Installation Start: 26-Apr-04	Datum	Elevations on NAVD88	Elevations on Mean Lower Low Water	Elevations on Mean Sea Level
Station Home Page: 8658163, Wrightsville Beach, NC	MHHW	0.526	1.367	0.709
Station Datum Definitions: 8658163, Wrightsville Beach, NC	MHW	0.421	1.262	0.604
	MSL	-0.183	0.658	0
RSLR: N/A	MLW	-0.793	0.048	-0.61
	MLLW	-0.641	0	-0.658

Data Inventory APALACHICOLA, FL

DEMS

Note: Rows shaded in orange indicate no bathymetry, yellow shading indicates that the vertical datum was something other than NAVD 88 for bathymetry.

Source	Resolution	Tidal Datum	Relative Tidal Datum	Vertical Datum	Vertical Units	Date Published	Decade Represented	Notes
NOAA NCEI	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2019	2010s	Topobathy. Site mosaicked from 2 individual 15 minute x 15 minute rasters. Bathymetric and topographic data utilized for DEM creation originate from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE. DEMs are referenced vertically to NAVD 88 with vertical units in meters.
USGS NED	~3m - 1/9 arc second	N/A	N/A	NAVD88	Meters	2007-2010	2000s	Topography only. Site mosaicked from 9 individual 15 minute x 15 minute rasters. The NED is derived from diverse source data that are processed to a common coordinate system and unit of vertical measure.
NOAA NCEI CRM Vol. 2 (THREDDS)	~90m - 3 arc second	MLLW (EPSG: 5866)	MSL	NAVD88 for topography	Meters	2001	1990s	Topobathy. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW). Source topographic data were in NAVD 88. The differences between these datums are less than the vertical accuracy of the CRM, so you can assign MSL to the CRM if you like, just recognize that the elevation values may not be as accurate as you might like or need. Assume a vertical accuracy no better than 1 meter for any elevation values in the CRM.

DEM CORRECTIONS

Data for RTK surveys conducted by Michael Starek available for the following locations and dates:

Location	Date
Cat Point	05/26/21; 09/22/21
East Bay	05/27/21
Unit 4	03/18/16; 03/25/17; 07/11/18; 05/24/19; 09/22/20; 05/27/21

ECOLOGICAL DATA

See note below tables for details regarding the Apalachicola NERR Vegetation Survey

Location	Type	Date	Source
Apalachicola NERR	Point Salinity from SET	2020	Apalachicola NERR
Apalachicola NERR	Point Salinity	2021	http://cdmo.baruch.sc.edu/
Apalachicola NERR	Vegetation Survey	2015 - 2020	Florida DEP


"The objectives of the APA NERR monitoring program are consistent with the objectives outlined in the NERRS technical report on Long-term Monitoring of Estuarine Submersed and Emergent Vegetation (Moore 2013) with some adaptations specific to the APA NERR. Specifically, the objectives of the APA NERR emergent vegetation protocol are: 1. Quantify vegetation patterns and their change over time and space; 2. Be consistent with other monitoring programs used worldwide, especially those used at other NERRS; 3. Be consistently applicable over a wide range of estuarine sites and habitats, including mangrove forests and seagrass meadows; 4. Quantify relationships among the various edaphic factors and the processes that are regulating the patterns of distribution and abundance in marsh communities; 5. Provide detailed information that can supplement comprehensive remotely sensed mapping of vegetation communities and other NERRS System Wide Monitoring Program data collection, as well as NERRS/NOAA education, stewardship, and restoration efforts."

SEA LEVEL TRENDS

Nonlinear trends for Apalachicola, FL


Created using data from the Interagency Sea Level Rise Scenario Tool

https://sealevel.nasa.gov/task-force-scenario-tool?psmsl_id=1193



Relative Sea Level (linear) trends for Apalachicola, FL

From NOAA: https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8728690



25-year linear rate of SLR from 1995-2020 for Apalachicola, FL

5.84 mm/yr with a 95% CI of 1.65

GAUGE DATA

Present Installation Start: 22-Sep-89	Datum	Elevations on NAVD88	Elevations on Mean Lower Low Water	Elevations on Mean Sea Level
Station Home Page: 8728690, Apalachicola, FL	MHHW	0.26	0.492	0.215
Station Datum Definitions: 8728690, Apalachicola, FL	MHW	0.228	0.46	0.183
RSLR: 8728690, Apalachicola, FL	MSL	-0.045	0.277	0
	MLW	-0.11	0.122	-0.155
	MLLW	-0.232	0	-0.277

Data Inventory GRAND BAY, MS

DEMS

Note: Rows shaded in orange indicate no bathymetry, yellow shading indicates that the vertical datum was something other than NAVD 88 for bathymetry.

Source	Resolution	Tidal Datum	Relative Tidal Datum	Vertical Datum	Vertical Units	Date Published	Decade Represented	Notes
MARIS/MDEQ	Varies	N/A	N/A	NAVD88	Meters (Converted)	2019	2010s	Mosaic of Lidar Projects spanning 10 years in Mississippi. State-wide Mosaic reports all elevations in US Feet. Original elevations were mixed between Meters and Feet.
USACE NCMP	1m	N/A	N/A	NAVD88	Meters	2018	2010s	Rasterized topobathy lidar elevations generated from data collected by the Coastal Zone Mapping and Imaging Lidar (CZMIL) system. CZMIL integrates a lidar sensor with simultaneous topographic and bathymetric capabilities, a digital camera and a hyperspectral imager on a single remote sensing platform for use in coastal mapping and charting activities.
USGS NED	~3m 1/9 arc second	N/A	N/A	NAVD88	Meters	2012	2000s	NED is derived from diverse source data that are processed to a common coordinate system and unit of vertical measure. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters digital hydrographic surveys which cover the central Gulf of Mexico region including Grand Bay.
USGS NED	~3m 1/9 arc second	N/A	N/A	NAVD88		2002	1990s	

DEM CORRECTIONS

Data for RTK surveys conducted by Jonathan Pitchford available for the following location and dates:

Location	Date
Grand Bay, MS	2013 and 2018
Available for download at Digital Coast: https://bit.ly/3JAXj1T	

ECOLOGICAL DATA


All NERR Point Salinity data are available from <http://grandbaynerr.org/data-downloads/>

Location	Type	Date	Source
Grand Bay, MS	Point Salinity and Dominant Vegetation	2020	Jonathan Pitchford
Bayou Cumbest	Point Salinity	2004 - present	NERR
Bayou Heron	Point Salinity	2004 - present	NERR
Bangs Lake	Point Salinity	2004 - present	NERR
Crooked Bayou	Point Salinity	2004 - 2005	NERR
Point Aux Chenes Bay	Point Salinity	2005 - present	NERR

SEA LEVEL TRENDS


Nonlinear trends for Dauphin Island, AL

Created using data from the Interagency Sea Level Rise Scenario Tool
https://sealevel.nasa.gov/task-force-scenario-tool?psmsl_id=1156



Relative Sea Level (linear) trends for Dauphin Island, AL

From NOAA: https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8735180



25-year linear rate of SLR from 1995-2020 for Dauphin Island, AL

7.01 mm/yr with a 95% CI of 1.84

GAUGE DATA

Present Installation Start: 25-May-10	Datum	Elevations on NAVD88	Elevations on Mean Lower Low Water	Elevations on Mean Sea Level
Station Home Page: 8740166, Grand Bay NERR, Mississippi Sound MS	MHHW	0.302	0.485	0.249
Station Datum Definitions: 8740166, Grand Bay NERR, Mississippi Sound MS	MHW	0.273	0.456	0.220
	MSL	-0.053	0.236	0
	MLW	-0.144	0.039	-0.197
RSLR: N/A	MLLW	-0.183	0	-0.236


B4: Map Data Exploration Activity Sheet

Map Data Exploration Activity Sheet





Link to map: <https://arcg.is/0CL9D90>

1. How does the NWI land cover data differ from the CCAP land cover data in Apalachicola?
 2. How many SETs show a negative elevation trend in Apalachicola?
 3. In Apalachicola, there are a number of clusters of RTK readings near Cat Point Rd. What is the elevation of the RTK point named "Cat Point – GCP18"?
 4. How do elevation trends (i.e., SET trends) in Alabama & Mississippi compare to those in Apalachicola?
 5. What appears to be the most common CCAP land cover type, or types, in Alabama & Mississippi?
 6. What is the name of the northernmost SET in North Carolina?
 7. Which of the three locations had the most SET sites?
 8. When looking at the DEM information layer, what colors represent the highest and lowest elevations?
 9. In the 2010s DEM for North Carolina, what does the elevation appear to be on Harker's Island?
 10. Does the DEM data differ across time in each location?
 11. For the DEM data available in Alabama & Mississippi, what year(s) of DEM data are available?
-

Appendix C: Data Exploration Web Application Screenshots




Stakeholder Data Investigation Tool



This web mapping application was developed to inform a retrospective marsh modeling workshop occurring in Spring 2022. It leverages a combination of digital elevation models (DEMs), digitized tidal datums, land cover data, and point locations for SET and RTK resources.

Data is presented for several NERR system locations in Mississippi, Florida, and North Carolina. Pages and data resources are represented in decadal bins split by state with source data assigned to a representative decade.

North Carolina



Navigate Here

Sites
North Carolina sites include the Currituck Banks, Rachel Carson, and Masonboro Island NERRs.


Elevation
2010s representative DEMs for all locations were sourced from the NOAA NCEI published between 2018 and 2019 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources, including the NOAA OCS, NOAA NGS, NOAA OCM, USGS, and USACE.

2000s representative DEMs for all locations were sourced from the USGS NED published in 2003 at a resolution of 1/9 arc seconds (approximated to 3 meters). This resource references the NAVD88 vertical datum with units in meters. The Coastal Relief Model (CRM) was used as the bathymetry resource for both the Rachel Carson and Currituck Banks sites. Best available bathymetric data were selected with a GIS query procedure that applied spatial and temporal filters to the 122 digital hydrographic surveys, dating from 1870 to 2005, which cover the North Carolina region.





1990s representative DEMs for all locations were sourced from the NOAA NCEI CRM Volume 2 published in 1998 at a resolution of 3 arc seconds (approximated to 90 meters). Bathymetric and topographic data utilized in the creation of this resource originated from a variety of sources. The vertical datum for the source bathymetric data was generally mean lower low water (MLLW) with source topography in NAVD88.

Tidal Datum Gauges

- Currituck Banks - 8651370, Duck, NC
- Rachel Carson - 8656483, Beaufort, Duke Marine Lab, NC
- Masonboro Island - 8658163, Wrightsville Beach, NC



Stakeholder Data Investigation Tool



Grand Bay

2000s

Grand Bay NERR Boundary

Grand Bay RTK

Grand Bay SET

Elevation Trend

Positive

2000s Representative Tidal Contours

Tidal Datum


- MLLW
- MLW
- MSL
- MHW
- MHHW

National Wetlands Inventory - Grand Bay

Wetland Type

Select a Different Decade

'90s '00s



3 km
3 mi

Esri, Maxar, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS


Powered by Esri

View a Different Location

Alabama & Mississippi


North Carolina

Florida



Appendix D: Workshop Flip Charts

D1: Workshop Marina & Ground Rules



D2: Managers Want Modelers to Know

Managers Want Models That:

- Are easy to understand
- Are specific to their managed lands
- Include outputs for various timesteps and SLR scenarios
- Are transparent about the input data

- Are clear about what any uncertainty means
- Can analyze management options

D3: Things for Managers to Know

Things for Managers to Know

1. Age of NWI data
2. Clarify diff. btwn time-steps & updates
3. Stage of development = generalities \rightarrow tipping points by decade not year
4. "elevation capital"
5. Managers = tuners
6. vulnerability vs "the future"
7. Models to understand process & vulns

D4: Data Discussion

Data Discussion

- NWI classifications missing
- glad to SETs, historic DEM
- concerns w/ getting historic data of NWI / CCAP \rightarrow Δ in classification
- SETs aren't randomly distributed \downarrow reanalyze original imagery
- WQ that includes Salinity & Sediments
- concerns w/ DEM resolution
- need built infra data
- dated sediment cores
- data trends ^{points} on sea level
? Stream gauges
- layers that track Δ in habitat
- consider ~~to~~ developing standardized data out of multiple datasets
- different sensitivities to DEMs in different systems
(... 10 m)

D5: Details of the Retrospective

Veg Data

- up to date veg data
- biomass; elevation = mean
- species mixture } CPRA
- SET accretion
- retrospective needs
- Veg Classes
 - High/Low tidal = constant sp mix
 - good for SET but diff cor
- Dominant species } warmer

Veg Data

- calibrating based on current imagery
- ~~inputs are~~
- species response comes from today assume in initial condition → can also do for a community
- Not CCAP b/c of lack of differentiation
- Aim for = H, L, Low medium
- need to define H, L, Low

DEM Data

- SLAMM
- Titus
- needs
- adapt
- all start
- merge
- Coris
- DEM current
- dense
- bet DEMs from SET data
- Time Steps
- 5 years - elevation
- compare
- available
- site

DEM Data

- SLAMM generates LiDAR
- Titus paper (Titus finding)
- needs to be LiDAR data
- post 09/05 hurricane
- 10s in other areas
- adapt historic data for all starting inputs?
- Coris → subtract diff from
- DEM data
- dense adjusted DEM from bet DEMs multiple spatial cores
- SET data, High/Low/Low/Low/Low
- Time Steps
- 5 years - compare point elevations
- compare landscape based on available lidar & veg data
- site variable

SLR data

- SLAMM assumes longer
- annual datapoints
- can do a steady state
- plan do both
- real data & smooth curve
- use best fit curve per station

Location Data

- a site historic
- Grand Bay
- lots of data
- Elevation
- NC
- Plum Island

SLR data

- SLAMM assumes longer time steps
- annual datapoints
- can do a steady state
- plan do both
- real data & smooth curve
- use best fit curve per station

Location Data

- a site historic
- Grand Bay
- lots of data
- Elevation
- NC
- Plum Island

Other Stuff

- Uncertainty
 - in how it's run
 - characterizing for end-users
 - structural vs parameter uncertainty
- Standardize
- an opportunity

SLAMM, WARMER
LA, MEM → range of conditions

D6: Scoping Question Discussion

Scoping Disc.

- teasing apart SLR & estuary characterization? (e.g. sal)
 - ↳ ideally understand all the variables are likely to influence output.
- challenges in the past
 - * signal of SLR
 - * amount of Δ
 - * by other factors (e.g. tagging)
- ② estuary factor suggestion
 - * minimize other factors of Δ besides SLR
- need to account for land management that has already occurred
 - * think about how to consider this * idios
- estuary factor: different hydrologic & geomorphic settings
 - * take lessons from our

Other
Uncertainty

D7: Next Steps

Data Wrangling

	Time	How?	Who?
Get NWI data: <small>(locations & bear spots on waterways)</small>	2 mo	no	Chris, me, Noah
Classify NWI data:	6-8 mo	yes	Jessica, Eric
Get NDEM data: <small>(bear locations)</small>	2-4 mo	no	Marion, Eric, Christine
Pick LA location:	2 mo	no	Eric, Christine
Existing cores: <small>- determine cores need</small>	4 mo	no	Tom - GB, Eric - LA, Maria - PI, Jim
Get SLR data:	ASAP	Jim, Eric, Trevor	
Get PIT data:	8 mo	yes	Tanner
Build Frank's NDEM:	1.5 years	yes	Tanner, GB, Eric - LA, Marion - PI
Get specific model costs	2 years	yes	hire out
	1 mo	No	Trevor, Eric, Marion, Jim

Post-it

30

23 INPT x 30 INPT
63.5 mm x 76.2 mm
52-550 FFP95 (0.48 µm)

3M

email updates → quarterly

meeting checkins

semi-annual * as needed

